

VOLUME 13 NO. 2

DECEMBER 2016

ISSN 1675-7009

SCIENTIFIC RESEARCH JOURNAL

Institute of Research Management & Innovation (IRMI)

Analysis of *Pandanus* Extracts by Reversed Phase High Performance Liquid Chromatography

Hannis Fadzillah Mohsin, Ratni Suriyani Jalal, Ibtisam Abdul Wahab, Abdul Rasyid Zulkifli & Muhammad Faxrurrazy Sadiran

Vitex Species: Review on Phytochemistry and Pouch Design for Nutritional Benefits

Nur Rejai Salmah Abdul Hakeem, Nuruljannah Md Yusof, Aisyah Hasyila Jahidin, Mizaton Hazizul Hasan, Hannis Fadzillah Mohsin & Ibtisam Abdul Wahab

Effect of Solar Radiation to the Building Materials Properties: A Review

Raihana Mohamad Hata, Rohana Hassan, Fadzil Arshad & Haslin Idayu

Adopting Industrialisation in the Construction Environment: Meeting the Requirements of Sustainability

Muhamad Faiz Musa, Mohd Reeza Yusof, Noor Sahidah Samsudin, & Faridah Muhamad Halil

Physical and Mechanical Properties of *Tenun Pahang* Fabrics using Alternative Yarns

Engku Liyana Zafirah Engku Mohd Suhaimi, Jamil Salleh, Suzaini Abd Ghani, Mohamad Faizul Yahya & Mohd Rozi Ahmad

SCIENTIFIC RESEARCH JOURNAL

Chief Editor

Hamidah Mohd Saman
Universiti Teknologi MARA, Malaysia

International Editors

David Shallcross, University of Melbourne, Australia
Ichsan Setya Putra, Bandung Institute of Technology, Indonesia
K. Ito, Chiba University, Japan
Luciano Boglione, University of Massachusetts Lowell, USA
Vasudeo Zambare, South Dakota School of Mines and Technology, USA

Editorial Board

Halila Jasmani, Universiti Teknologi MARA, Malaysia
Kartini Kamaruddin, Universiti Teknologi MARA, Malaysia
Tan Huey Ling, Universiti Teknologi MARA, Malaysia
Mohd Zamin Jumaat, University of Malaya, Malaysia
Norashikin Saim, Universiti Teknologi MARA, Malaysia
Noriham Abdullah, Universiti Teknologi MARA, Malaysia
Saadiah Yahya, Universiti Teknologi MARA, Malaysia
Norizzah Abdul Rashid, Universiti Teknologi MARA, Malaysia
Zahrah Ahmad, University of Malaya, Malaysia
Zulkiflee Abdul Latif, Universiti Teknologi MARA, Malaysia
Zulhabri Ismail, Universiti Teknologi MARA, Malaysia
Ahmad Zafir Romli, Universiti Teknologi MARA, Malaysia
David Valiyappan Natarajan, Universiti Teknologi MARA, Malaysia
Fazlena Hamzah, Universiti Teknologi MARA, Malaysia
Nor Ashikin Mohamed Noor Khan, Universiti Teknologi MARA, Malaysia
Sabarinah Sheikh Ahmad, Universiti Teknologi MARA, Malaysia
Ismail Musirin, Universiti Teknologi MARA, Malaysia
Norhati Ibrahim, Universiti Teknologi MARA, Malaysia
Kalavathy Ramasamy, Universiti Teknologi MARA, Malaysia
Ahmad Taufek Abdul Rahman, Universiti Teknologi MARA, Malaysia

Journal Administrators

Yazmin Sahol Hamid, Universiti Teknologi MARA, Malaysia
Khairul Nurudin Ahnaf Khaini, Universiti Teknologi MARA, Malaysia

© UiTM Press, UiTM 2016

All rights reserved. No part of this publication may be reproduced, copied, stored in any retrieval system or transmitted in any form or by any means; electronic, mechanical, photocopying, recording or otherwise; without prior permission in writing from the Director of UiTM Press, Universiti Teknologi MARA, 40450 Shah Alam, Selangor Darul Ehsan, Malaysia. E-mail: penerbit@salam.uitm.edu.my.

The views, opinions and technical recommendations expressed by the contributors and authors are entirely their own and do not necessarily reflect the views of the editors, the publisher and the university.

SCIENTIFIC RESEARCH JOURNAL

Institute of Research Management & Innovation (IRMI)

Vol. 13, No. 2	December 2016	ISSN 1675-7009
1.	Analysis of <i>Pandanus</i> Extracts by Reversed Phase High Performance Liquid Chromatography <i>Hannis Fadzillah Mohsin</i> <i>Ratni Suriyani Jalal</i> <i>Ibtisam Abdul Wahab</i> <i>Abdul Rasyid Zulkifli</i> <i>Muhammad Faxrurrazy Sadiran</i>	1
2.	<i>Vitex</i> Species: Review on Phytochemistry and Pouch Design for Nutritional Benefits <i>Nur Rejai Salmah Abdul Hakeem</i> <i>Nuruljannah Md Yusof</i> <i>Aisyah Hasyila Jahidin</i> <i>Mizaton Hazizul Hasan</i> <i>Hannis Fadzillah Mohsin</i> <i>Ibtisam Abdul Wahab</i>	15
3.	Effect of Solar Radiation to the Building Materials Properties: A Review <i>Raihana Mohamad Hata</i> <i>Rohana Hassan</i> <i>Fadzil Arshad</i> <i>Haslin Idayu</i>	29

4. **Adopting Industrialisation in the Construction Environment: Meeting the Requirements of Sustainability** 45
Muhamad Faiz Musa
Mohd Reeza Yusof
Noor Sahidah Samsudin
Faridah Muhamad Halil
5. **Physical and Mechanical Properties of *Tenun Pahang* Fabrics using Alternative Yarns** 67
Engku Liyana Zafirah Engku Mohd Suhaimi
Jamil Salleh
Suzaini Abd Ghani
Mohamad Faizul Yahya
Mohd Rozi Ahmad

***Vitex* Species: Review on Phytochemistry and Pouch Design for Nutritional Benefits**

Nur Rejai Salmah Abdul Hakeem, Nuruljannah Md Yusof,
Aisyah Hasyila Jahidin, Mizaton Hazizul Hasan, Hannis Fadzillah Mohsin¹
and Ibtisam Abdul Wahab

*Department of Pharmacology & Chemistry, Faculty of Pharmacy,
Universiti Teknologi MARA Selangor, Puncak Alam Campus,
42300 Bandar Puncak Alam,
Selangor Darul Ehsan*

¹E-mail: hannisfadzillah@gmail.com

ABSTRACT

Vitex trifolia or locally known as *lemuni*, is well documented for its medicinal properties including anticancer, antibacterial and wound healing. For years, the leaves and flowers were consumed for anti-aging and general well-being. It is also commonly served as nutraceuticals to post-partum women during their confinement period. Previous study showed that *V. trifolia* possesses high antioxidant properties that could prevent oxidative stress related diseases such as cancer, atherosclerosis, diabetes and many more. From the literature, *V. trifolia* leaves are used as poultice for rheumatic pains, sprains, antihelmintics and inflammations. Meanwhile, the fruits are utilised in amenorrhea. The roots, on the other hand, are used for the treatment of cough and fever. The plant is known to possess pharmacologically-active constituents such as the essential oil, halimane-type diterpenes, flavonoids, triterpenes, lignans, iridoids and ecdysteroids. Review papers were consistently published to justify the non-exhaustive biological exploration of this genus. Nevertheless, only 11 per cent of the total *Vitex* species were phytochemically examined. Apart from being eaten raw or blanched, *Vitex* is also taken as a food colourant in preparing *lemuni* rice. The process of making *lemuni* rice is quite tedious. First, the leaves and flowers are picked and washed. Both parts are then ground with water and sieved to give a bluish extract which is then mixed with rice and boiled to cook. Hence, the *Vitex* pouch is introduced to provide an easy way to prepare *lemuni* rice. This packaging is designed to enclose *Vitex* samples in a pouch that simply can be placed together with rice and boiled to cook.

Such instant preparation will hopefully promote the consumption of lemuni rice among busy locals. Future plans include international marketing and production of other Vitex nutritional products.

Keywords: *nutrition, phytochemistry, pouch, spectroscopy, vitex*

INTRODUCTION

Vitex species, locally known as lemuni, contain a variety of potentially bioactive molecules, such as iridoids, flavonoids, diterpenoids, derivatives and phytosteroids. These plants were used in traditional medicine to cure a wide range of ailments, for example, depression, allergy, wounds, diarrhoea and also gastrointestinal diseases [1]. A number of biological activities of this species were reported. Besides the antioxidant and antileptic activities of *V. agnus-castus*, *V. trifolia* also possesses several pharmacological properties such as hepatoprotective, wound healing activity and reducing allergy disease [2]. Meanwhile, *V. negundo* was proven to have analgesic and anti-inflammatory actions. In India, *V. penduncularis* and *V. pinnata* were found to have insecticidal properties [2]. Among 250 species in the *Vitex* genus, *V. trifolia* is one of the very well-known species and is commonly being used in *Vitex* studies. Based on research, this species is believed to have various therapeutic benefits. The plant is known to possess biologically-active constituents such as the essential oil, halimane-type diterpenes, flavonoids, triterpenes, lignans, iridoides and ecdysteroids [1]. It is widely used in India as the remedies in their Ayurvedic and Unani systems of medicines [3].

Most parts of *Vitex* plants (the leaves, fruits, roots and stems) are known to have many medicinal values. However, the specific chemical constituents that present in each part are still unclear. The importance of understanding which chemical constituents in each plant part can help to determine the compounds that are responsible for the therapeutic benefits. Thus, by extracting these *Vitex* species based on individual parts, the detailed information on the chemical compound could be understood. In Asian countries, *V. trifolia* leaves are consumed to improve memory, remove bad taste in mouth, cure fever and as a pain reliever [4]. The leaves are used internally or externally in baths to cure Ciguatera fish poisoning-related pruritus. Besides its antipyretic activity, it can also increase body weight

and being reported to have anti-tumour activity [2]. It can also being used as nematocidal and antihelminthic agents. The Asians choose *V. trifolia* as their remedy due to its ability to cure rheumatic pains, sprains, leucoderma and bronchitis. It is also utilised for hair growth and eye vision improvement [5]. By monitoring its effects on the modulation of cytokines, mediators of inflammation, the leaves part is believed to have anti-inflammatory property [2].

Regarding the other plant parts, the roots are used to treat febrifuge, cough and fever. The stems of *V. trifolia* was proven to be very toxic against cultures of several cell lines [2]. The aerial parts of this plant are useful in diabetes treatment. Meanwhile, the leaf extract is anti-cancerous. The fruits, in the other hand, are good for amenorrhoea. When the flowers of this *Vitex* species were mixed with honey, they were used in fevers, accompanied with severe thirst [4]. Besides *V. trifolia*, *V. negundo* is also a well-known species. In addition to this species being easily found and collected, *V. negundo* also possesses a wide variety of compounds. Although it does not have therapeutic aids as much as *V. trifolia*, studies of *V. negundo* have afforded numbers of phytochemicals, such as volatile oils, lignans, flavonoids, terpenes, and steroids [2]. The plant analysis showed that its leaves contains alkaloid (nishundine), luteolin-7-glucoside, vitexicarpin 1 (or casticin, 3',5-dihydroxy-3,4',6,7-tetramethoxyflavone, Figure 1), iridoid glycosides, an essential oil and other constituents like vitamin C, carotene, gluco-nonital, benzoic acid, β -sitosterol and C-glycoside. The seeds contain hydrocarbons, β -sitosterol, benzoic acid and phthalic acid, diterpenoids, flavonoids, artemetin and triterpenoids. Meanwhile, the bark contains fatty acids, valinic acid, p-hydroxybenzoic acid and luteolin [6].




Figure 1: The chemical structure of vitexicarpin 1

Studies demonstrated that vitexicarpin 1 could be isolated from the flowers of *Vitex rotundifolia* (*Vitex rotundifolia* Linne fil.). It has long been used as an anti-inflammatory herb in traditional Chinese medicine [7].

A number of recent pharmacological reports indicated that 1 inhibits the proliferation of lymphocytes and thus, inhibit the growth of cancer cells in various cell lines [7]. Previously, a study on the leaves of *V. trifolia* yielded three compounds, viteosin A, 1 and vitetrifolin E. Vitexicarpin 1 was the most active among the three compounds. The mechanism of activity seems to be non-competitive antagonism to histamine and stabilisation of mast cells membrane function. The leaves extract gave an essential oil. Meanwhile, the fruits contain an acid resin, an astringent organic acid, malic acid, traces of an alkaloid and colouring matter [8].

In this study, the focus is given in the separation and identification of the chemical composition in the leaves extract of a *Vitex* species by using standard chromatographic technique. Apart from the phytochemical analysis, the *Vitex* pouch is also introduced, in order to provide an alternative for the preparation of lemuni rice. Traditionally, *Vitex* is eaten raw or blanched. It is also consumed as a food colourant in preparing lemuni rice. Nevertheless, the process of making lemuni rice is quite tedious. The leaves and flowers were picked and washed, before they were ground with water and sieved to give a bluish extract. This colorant is then mixed with rice and boiled to cook. Hence, the *Vitex* pouch is proposed to provide convenient means to prepare lemuni rice.

METHODOLOGY

Raw material

The crude drug, which is the flowering stems of *V. trifolia* was collected from Klang, Selangor Darul Ehsan, Malaysia, in July 2011. A voucher specimen (no.0711VKLG) was deposited in Faculty of Pharmacy, Universiti Teknologi MARA, Puncak Alam, Selangor Darul Ehsan, Malaysia. The plant material was later divided into leaves, fruits and twig parts.

Chemicals

The chemicals that were used include the solvents; chloroform (CHCl_3), ethanol (EtOH) and methanol (MeOH). Ceric sulphate was used as the spray reagent to visualise ultraviolet (UV) inactive compound.

Extraction, Isolation and Purification of the *Vitex* Extract

The plants parts were cut before they were extracted. The leaves were introduced to EtOH extraction, silica column and preparative thin layer chromatography (TLC). Plates coated with silica gel (silica 60 F254) was used as the stationary phase. A pure compound 1 was purified and subjected to Nuclear Magnetic Resonance (NMR) spectroscopy (500 MHz, CD_3OD).

Designing *Vitex* Pouch for Fresh Herbal Packaging

Vitex is cherished in local rice preparation; however, this herb is seen as having relatively low commercial value in food industry. Therefore, there is a call to incorporate *Vitex* natural materials into new food products. For travellers or tourists, the difficulty of getting *Vitex* fresh material could be minimised with the provision of a novel form of this crop. Therefore, a packaging is designed to consist of ten grams of *Vitex*'s dried, shredded leaves, flowers and stem in a pouch that simply could be placed together with rice and boiled to cook (Figure 2). A pouch (11.0 cm x 10.5 cm) was utilised to enclose the powdered *Vitex* samples. For presentation purposes, two smaller pouches (9.5 cm x 7 cm) could be placed in a zip bag, having the dimension of 14 cm x 10 cm. An example of a complete product could consist of an outer case or box, the above pouch to dispense *Vitex*'s nutrient into the rice and a packet of rice, for example, dedicated for two travelling persons. Appropriate labelling could be designed for the boxes or packaging, in order to attract customers. In addition to *Vitex* pouch, other herbs such as the fragrant *Pandanus* leaves, a few cinnamon stick, clove, star anise and low-fat coconut milk could be added. Such instant preparation will hopefully promote the consumption of lemuni rice among busy locals. The shelf life of 12 months could be expected for the above food materials.




Figure 2: An example of the *Vitex* pouch and the dried leaves (source by author)

A preliminary sensory evaluation of the lemuni rice sample was carried out, four hours after its preparation [9]. A panel was randomly selected among the university staff for the evaluation. Nevertheless, a standard scale [10] was not utilised to evaluate the appearance, aroma, taste, mouth feel and acceptability of the sample. None of the necessary precautions were taken to prevent carrying over the taste perception. This evaluation is only meant as a general introduction to the *Vitex* product. Therefore, a more standardised sensory assessment will be performed in near future.

RESULT AND DISCUSSION

Isolation and Purification of the *Vitex* Extract

The samples were extracted and monitored via TLC analysis, in order to obtain their chromatographic profiles. A preliminary spectroscopic evaluation of an iridoid glycoside **2** is shown in the $^1\text{H-NMR}$ spectrum (500 MHz, CD_3OD , Figure 3). The signals at were assignable to aromatic protons, originated from a p-hydroxybenzoyl group (δH 6.8 - 8.0 ppm), three olefinic protons (δH 5.0 - 6.5 ppm), aliphatic protons and glucose residue (δH 3.2 - 5.0 ppm), another two aliphatic protons (δH 2.7 - 3.0 ppm), and finally $-\text{CH}_2$ protons, possibly adjacent to a carbonyl group; $-\text{CH}_2-\text{C}=\text{O}$ at δH 1.91 ppm.

The resonances attributed to a p-hydroxybenzoyl group [δ H 6.85 and 7.93, respectively H-2' (2H) and H-3' (AB system, doublets, $J = 7.5$ Hz) was observed. This characteristic moiety was frequently encountered in *Vitex* species. The acyl group was attached to C-10, and the C-10 methylene protons' resonances were found to be shifted downfield at δ H 1.91 (singlet, 2H). The signals of one doublet at δ H 6.37 ($J = 5.5$ Hz) and one doublets at δ H 5.14 ($J = 3.5$ and 5.0 Hz) were due to the olefinic protons at C-3 and C-4, respectively. The proton at C-4 was believed to couple to a proton at δ H 2.72 ppm (1H, multiplet, H-5), which could couple to a proton at δ H 4.49 (1H, multiplet, H-6). The chemical shift of H-6 suggested the presence of one hydroxyl group at C-6. The proton at C-6 might also correlate to an olefinic proton at δ H 5.82 (1H, broad singlet, H-7). In addition, the proton at C-5 could correspond with a proton at δ H 3.01 (1H, triplet, H-9).




Figure 3: ^1H -NMR (500 MHz, CD_3OD , δ H 0 - 9 ppm) spectrum of the suggested iridoid glycoside 1

From these findings, the data for compound 2 was recommended to be similar to those for agnuside, $\text{C}_{22}\text{H}_{26}\text{O}_{11}$ (Figure 4). This suggestion was also supported by comparison of the ^1H -NMR data [11-15]. The IUPAC name of 2 was established as [5-hydroxy-1-[3,4,5-trihydroxy-6-(hydroxymethyl)-oxan-2-yl]oxy-1,4a,5,7a-tetrahydrocyclopenta[c]pyran-7-yl]methyl 4-hydroxy-benzoate. Hence, compound 2, was proposed as a known composition of *V. trifolia* and it is commercially available [16]. The occurrences of 2 were also reported in other *Vitex*, including *V. cannabifolia*, *V. rotundifolia*, *V. cymosa*, *V. agnus-castus* and *V. negundo*. This compound could be seen to compose of aucubin 3 (Figure 5) and p-hydroxybenzoic

acid, another natural component of *Vitex* species. The presence of the both compounds 2 and 3 are quite common in almost *Vitex* species. Hence, the above compounds may be considered as chemotaxonomic markers of the genus *Vitex* [17]. The quantification of 2 via high performance liquid chromatography (HPLC) proposed that the leaves contained relatively high amount of 2. Therefore, the bark part should not be disrupted, thus setting the efforts in conserving the plant [16].




Figure 4: Compound 2 was proposed as agnuside, a known composition of *V. trifolia*




Figure 5: The chemical structure of aucubin 3

Panel Suggestion of the *Vitex* Pouch and Lemuni Rice

The plant materials were stored to dry, naturally, before they were used for the pouch. Bigger, industrial scale could apply an oven drying, instead. Other methods of preparation would include; either freeze drying of *Vitex*'s aqueous extract; or by using liquid nitrogen. However, these two

approaches would not assure the freshness of the colourant and its antifungal properties. On the other hand, the pouch design could be considered as a type of traditional packaging, due to its basis of prior drying of the plant materials as the primary preservation method [18]. A more modern approach would be to convert the concentrated *Vitex* extract into cubes, as seen for chicken/beef stock cubes, possibly with added chemical preservatives or artificial colours. The cubes would later be broken up or dissolved with boiling water while the lemuni rice is cooked.

The introductory assessment of the pouch and lemuni rice were carried out by a team of 13 members. It was apparent that the rice taste is contributed by the added spice in the lemuni rice, such as cinnamon and star anise. The volatile and non-volatile components of *Vitex* and the spices give the overall flavour and aroma. Specifically, the volatile composition of the *Vitex* extractives, which is also referred as the essential oils, could benefit the consumer, since the essential oils of *V. trifolia* was shown to possess insecticidal properties [2]. In addition, the dark-blue characteristic colouring of the rice provide a distinct, vivid impact to the rice, as compared to the plain, white rice. The panel also suggested that the unique taste and rich aroma of *Vitex* leaves was well blended with the rice. It was summarised that the panels altogether liked the lemuni rice, prepared with the designed pouch.

CONCLUSION

Iridoids and terpenoids are frequently discovered from this traditional plant. Extensive experiments, including two-dimensional NMR, will be carried out to fully characterise this plant's natural metabolite. Review papers were consistently published to justify the non-exhaustive exploration of this genus, both phytochemically and biologically [19-21]. On another note, such instant lemuni rice preparation will hopefully promote *Vitex* consumption among busy locals. The design and concept of the *Vitex* pouch could be realised and disseminated to the public. With this proposal, the herbal product could be introduced and acquired for a quick, convenient and effective preparation of lemuni rice. The temperature range of about 80-100°C will occur during the rice cooking process, thereby infusing

the *Vitex*'s nutrients into the rice. Generally, most food products can be constantly reformulated, therefore, the natural ingredients and its nutrition content may change. Therefore, further invitation should be extended to food engineers, biotechnologists and nutritionists, in order to investigate this point. The future plans would include international marketing and production of other *Vitex* nutritional products.

ACKNOWLEDGEMENT

The authors would like to thank UiTM for the undergraduate research opportunity. The authors would also like to acknowledge Mr. Mohd Syukri Baharudin and Miss Nawal Huda Rafeek Ahmad @ Mohamed Towfik for the technical support.

REFERENCES

- [1] M. A. Mohamed, A. M. Abdou, M. M. Hamed and A. M. Saad, 2012. Characterization of bioactive phytochemical from the leaves of *Vitex trifolia*, *International Journal of Pharmaceutical Applications*, Vol. 3(4), pp. 419-428.
- [2] A. K. Meena, U. S. Niranjana, M. M. Rao, M. M. Padhi and R. Babu, 2011. A review of the important chemical constituents and medicinal uses of *Vitex* genus, *Asian Journal of Traditional Medicines*, Vol. 6(2), pp. 54-60.
- [3] M. K. Aswar, A. A. Bidkar, K. N. Gujar and T. G. Athawale, 2012. Anxiolytic Like Effects of Leaves Extract of *Vitex negundo* (L) - (Fam:Verbaseae) in Elevated Plus Maze Test, *Journal of Natural Remedies*, Vol. 12(2), pp. 141-150.
- [4] N. Tiwari, J. Thakur, D. Saikia and M. M. Gupta, 2013. Antitubercular diterpenoids from *Vitex trifolia*, *Phytomedicine*, Vol. 20(7), pp. 605-610.
- [5] S. Thenmozhi, K. Vibha, M. Dhanalakshmi, K. Manjuladevi, S. Diwedi and U. Subasini, 2013. Evaluation of Anthelmintic Activity of *Vitex Trifolia* Linn Leaves against *Pheretima Posthuma*, *International Journal of Pharmaceutical & Biological Archives*, Vol. 4(5), pp. 878 – 880.
- [6] S. Das and L. Kanodia, 2013. Effect of Ethanolic Extract of Leaves of *Vitex negundo* L. on Acetic Acid Induced Colitis in Albino Rats, *Asian Journal of Pharmaceutical and Clinical Research*, Vol. 6(3), pp. 138-141.
- [7] F.-M. Meng, J. B. Yang, C. H. Yang, Y. Jiang, Y. F. Zhou, B. Yu and H. Yang, 2012. Vitexicarpin induces apoptosis in human prostate carcinoma PC-3 cells through G2/M phase arrest, *Asian Pacific Journal of Cancer Prevention*, Vol. 13(12), pp. 6369-6374.

- [8] W. G. Ko, T. H. Kang, S. J. Lee, N. Y. Kim, Y. C. Kim, D. H. Sohn and B. H. Lee, 2000. Polymethoxyflavonoids from *Vitex rotundifolia* inhibit proliferation by inducing apoptosis in human myeloid leukemia cells, *Food and Chemical Toxicology*, Vol. 38(10), pp. 861-865.
- [9] A. H. Jahidin, H. H. Mizaton and I. Abdul Wahab, 2015. *Vitex* Rice On-The-Go. Invention, Innovation & Design Exposition (IIDEX 2015), Universiti Teknologi MARA, 27th – 30th April 2015.
- [10] H. T. Lawless, 2013. *Laboratory Exercises for Sensory Evaluation*. New York: Springer.
- [11] T. Iwagawa, A. Nakahara, A. Miyauchi, H. Okamura and M. Nakatani, 1993. Constituents of the Leaves of *Vitex cannabifolia*, *Kagoshima Daigaku Rigakagu Kiyo, Suguku, Butsurigaku, Kagaku*, Vol. 26, pp. 57-61.
- [12] E. Okuyama, S. Fujimori, M. Yamazaki and T. Deyama, 1998. Pharmacologically active components of Vitis Fructus (*Vitex rotundifolia*). II. The components having analgesic effects, *Chemical Pharmaceutical Bulletin (Tokyo)*, Vol. 46(4), pp. 655-662.
- [13] T. C. Santos, J. Schripsema, F. D. Monache and S. G. Leitão, 2001. Iridoids from *Vitex cymosa*, *Journal of the Brazilian Chemical Society*, Vol. 12(6), pp. 763-766.
- [14] A. Kuruüzüm-Uz, K. Ströch, L. O. Demirezer and A. Zeeck, 2003. Glucosides from *Vitex agnus-castus*, *Phytochemistry*, Vol. 63(8), pp. 959-964.
- [15] F. Jamaludin, 2008. Chemical Constituents of *Vitex Negundo* and Evaluation of Their Anti-Inflammatory and Antioxidant Activities. Unpublished PhD's thesis, Universiti Putra Malaysia, Malaysia.

- [16] S. Shah, T. Dhanani and S. Kumar, 2013. Validated HPLC method for identification and quantification of p-hydroxy benzoic acid and agnuside in *Vitex negundo* and *Vitex trifolia*, *Journal of Pharmaceutical Analysis*, Vol. 3(6), pp. 500-508.
- [17] S. Ganapaty and K. N. Vidyadhar, 2005. Phytoconstituents and Biological Activities of *Vitex* - a Review, *Journal of Natural Remedies*, Vol. 5(2), pp. 75-95.
- [18] S. Y. Lee, S. J. Lee, D. S. Choi and Hur, S. J., 2015. Current topics in active and intelligent food packaging for preservation of fresh foods, *Journal of Science of Food Agriculture*. Vol. 95(14), pp. 2799 - 2810.
- [19] A. S. Vishwanathan and R. Basavaraju, 2010. A Review on *Vitex negundo* L. – A Medicinally Important Plant, *EJBS*. Vol. 3(1), pp. 30-42.
- [20] F. Basri, H. P. Sharma, S. Firdaus, P. Jain and A. Ranjan, 2014. A Review of Ethnomedicinal Plant-*Vitex negundo* Linn, *International Journal of Advanced Research*. Vol. 2(3), pp. 882-894.
- [21] C.J. Zheng, H.Q. Li, S.C. Ren, C.L. Xu, K. Rahman, L.P. Qin and Y.-H. Sun, 2015. Phytochemical and Pharmacological Profile of *Vitex negundo*, *Phytotherapy Research*. Vol. 29(5), pp. 633–647.

GUIDELINES FOR SUBMISSION OF ARTICLES

The Scientific Research Journal (SRJ) (ISSN: 1675-7009) is an international refereed journal, jointly published by the Institute of Research Management and Innovation (IRMI) and the UiTM Press, Universiti Teknologi MARA (UiTM), Malaysia. This journal was launched in the hope of stimulating quality research projects in science and technology related areas. Researchers are strongly encouraged to use this journal as a platform for publishing and disseminating their research findings to the members of academia and the community at large.

The SRJ publishes research papers that address significant issues in the fields of science and technology including life science, natural science, health science and physical science. It provides a balanced presentation of articles and solicits contributions from the fields of: Engineering, Computer Science, Mathematical Sciences, Energy, Environmental Science, Materials Science, Physics, Chemistry, Astronomy, Pharmacy, Medicine and Applied Sciences.

Emphasis is place on direct and clearly understood communication, originality and scholarly merit. Only original papers will be accepted and copyright of published papers will be vested in the publisher.

Submitted manuscripts should be written in English with production-quality figures. Manuscripts should be typed using Times New Roman (point 12), single-line spacing and wide margins (2.54 cm left, right, top and bottom). The manuscript should include the title of the paper; the name, address, contact numbers and email(s) of the correspondence author; a short abstract of between 200 and 300 words, which clearly summaries the paper with 4-5 keywords. Submission should be limited to a maximum of 20 typed pages.

Tables should be included within the text where appropriate and must be numbered consecutively with Arabic numerals and have titles that precede the table. Similarly, all figures must be numbered and a detailed caption should be provided below each figure. Figures should be embedded within the text where appropriate. Glossy photographs when required

should be scanned to a suitable resolution (1200 dpi), which enables quality reproduction.

References should be numbered in ascending order and cited within square brackets, e.g. [1], [3-5], in the main body of the text. References included at the end of the manuscript are to be in the order that they appear in the main body of text. Some entry samples are as follows:

- [1] A. B. Author, 2000. *Title of Book*, ABC Press, Kuala Lumpur, Malaysia.
- [2] C. D. Author and E. F. Author, 1999. Title of Paper, *Journal Name*, Vol. 10, pp. 32-45.
- [3] G. H. Author, 2006. Title of the conference paper, in Proceedings of the 2000 IEEE International Symposium on Circuits and Systems, Geneva, Switzerland, pp. 100-105

Manuscripts submitted to the journal will be initially screened by the Chief Editor, to determine appropriateness. Only those manuscripts considered of a sufficiently high standard will proceed to undergo a double blind review.

Upon completion, the manuscripts will be passed to an editorial board member for appraisal of their value. Additionally, they will be reviewed by an expert in that discipline.

For further information, please contact:

Professor Dr. Hamidah Mohd Saman

Chief Editor

Scientific Research Journal

Institute of Research Management and Innovation

Universiti Teknologi MARA

40450 Shah Alam Selangor,

MALAYSIA

E-mail: hamid929@salam.uitm.edu.my

Or e-mail SRJ's Secretariat: srj_editor@salam.uitm.edu.my/ khairul2950@salam.uitm.edu.my